

Cover Crops for Nitrogen Management



Eileen Kladviko, Agronomy Department, Purdue University

Created August 2011


URL: <http://www.ag.purdue.edu/agry/extension/Documents/CoverCropsNitrogen.pdf>

Cover crops can provide many benefits for soil and water quality and crop production. One of the important benefits of cover crops is their impact on soil nitrogen. Generally we would characterize cover crops as either a scavenger of soil N or a producer of soil N, although legumes will do both.

Scavenging N

Non-legume cover crops will scavenge or “trap” soil nitrate that would otherwise move out of the rootzone into tile drains or groundwater. Even with well-managed corn and soybean production, there is always some leaching of nitrate that originates either from residual fertilizer N or from the natural decomposition of soil organic matter. Our annual cropping systems are “leaky” because there are long fallow periods between crop maturity in September and the active growth of the next cash crop in May. Most of the net downward flow of water to the drains occurs precisely during this long fallow period, when there is nothing to take up the nitrate. Cover crops actively take up nitrate during a part of that fallow season, reducing the losses that occur to tile drains.

Corn-soybean system normally fallow from Oct – April.



Majority of drainflow and N-loads occur in fallow season



Illustration by Lou Jones/ E. Kladviko

Typical scavenger cover crops include grasses such as cereal rye and annual ryegrass, and brassicas such as oilseed radish. Legumes will also scavenge residual N before producing their own N, but because of their slower establishment they are not normally planted for the primary purpose of scavenging N. The amount of N scavenged depends on the amount of growth of the cover crop and the amount of N available in the soil. Fields with manure applications have greater available N and a greater need for a cover crop to trap the N. Typical corn/soybean systems without manure may have 10 to 40 lbs N/A trapped in the above-ground portion of the cover crop, while fields with manure may have 80 to 100 lbs/A or more. Cover crops also trap substantial amounts of N in their roots.

It is important to remember that non-legume cover crops are not producing or fixing N, they are simply scavenging or trapping N from the soil that would otherwise be subject to losses by leaching or denitrification.

Producing (or “fixing”) N

Legume cover crops fix atmospheric N for their own growth. The amount of N fixed depends on the amount of growth and the particular species. A challenge for inclusion of legumes in a corn-soybean system is the limited time available for growth and N-fixation. Legumes fit well after wheat, when not double-cropped, since there is more time for growth of the cover crop. Legumes must be inoculated before they will fix N.

Availability of the N scavenged or produced

How much of the N that was scavenged by a trap crop or fixed by a legume, becomes available for the next cover crop, and when? These are very challenging questions and there is no simple answer. The rate and extent of N release from the decomposing cover crop depends on many factors, including the age/stage of the plant when terminated, the nitrogen percent, carbon to nitrogen (C:N) ratio, plant composition, and weather (temperature and moisture). Plants in the vegetative stage have higher N concentrations than those in the reproductive stage and will decompose faster. The C:N ratio of the cover crop residues determine the general timing of release of N from the residues. With high C:N ratios (greater than 25:1), the decomposing cover crop residues will first immobilize N from the soil or recent fertilizer N additions. Only after some period of time will the N start to be released, or mineralized. When managing grass cover crops (rye, annual ryegrass, wheat), terminating the cover crop while it is still in the vegetative stage is generally the preferred strategy if growing a high N-user cash crop such as corn. If the cover crop gets into the reproductive phase before termination, there will be more N immobilization after termination and a good application of starter N fertilizer should be used.

Not all the N scavenged by the cover crop will be available the next season, so overall fertilizer N rate should not immediately be reduced by the amount of N kept out of the drain, for example. Some of the scavenged N goes towards building soil organic matter. After multiple years of cover crops and an increase in soil N content, there will likely be the possibility to reduce fertilizer N rates for a similar crop yield. This is an active area of experimentation by producers and researchers.

Legume cover crops and most of the grasses, while in the vegetative stage, have C:N ratios around 10:1 to 15:1. These decompose and release N rapidly after termination. Roughly about half of the N in the above-ground biomass will be available over a time window of around 1 to 2 months, depending on the weather. If a legume is terminated too early, some of the N may be lost before the cover crop can use it.

Summary

Cover crops can have important effects on soil and water quality by scavenging available N from the soil profile and recycling it. The recycled N helps build soil organic matter and some of it becomes available to succeeding crops. Legume cover crops fix atmospheric N, and upon termination, roughly half of the fixed N becomes available over a several month period. While there are many challenges remaining to fully integrate cover crops into nitrogen management strategies, they should be considered as an important part of increasing overall soil biological activity and soil quality.

© 2007-2011, Purdue University

It is the policy of the Purdue University that all persons shall have equal opportunity and access to its programs and facilities without regard to race, color, sex, religion, national origin, age, or disability. Purdue University is an Affirmative Action employer. This material may be available in alternative formats.